

# BINARY BROWN DWARFS IN THE GALACTIC HALO?

F. De Paolis<sup>1</sup>, G. Ingrosso<sup>2</sup>, Ph. Jetzer<sup>1</sup> and M. Roncadelli<sup>3</sup>

<sup>1</sup> *Paul Scherrer Institute, Laboratory for Astrophysics, CH-5232 Villigen PSI, and*

*Institute of Theoretical Physics, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland*

<sup>2</sup> *Dipartimento di Fisica, Università di Lecce, Via Arnesano, CP 193, 73100 Lecce, and*

*INFN, Sezione di Lecce, Via Arnesano, CP 193, 73100 Lecce, Italy*

<sup>3</sup> *INFN, Sezione di Pavia, Via Bassi 6, I-27100, Pavia, Italy*

## ABSTRACT

Microlensing events towards the Large Magellanic Cloud entail that a sizable fraction of dark matter is in the form of MACHOs (Massive Astrophysical Compact Halo Objects), presumably located in the halo of the Galaxy. Within the present uncertainties, brown dwarfs are a viable candidate for MACHOs. Various reasons strongly suggest that a large amount of MACHOs should actually consist of binary brown dwarfs. Yet, this circumstance looks in flat contradiction with the fact that MACHOs have been detected as unresolved objects so far. We show that such an apparent paradox does not exist within a model in which MACHOs are clumped into dark clusters along with cold molecular clouds, since dynamical friction on these clouds makes binary brown dwarfs very close. Moreover, we argue that future microlensing experiments with a more accurate photometric observation can resolve binary brown dwarfs.

**Key words:** Galaxy: halo - stars: brown dwarfs - dark matter - gravitational lensing

## 1 INTRODUCTION

As is well known (Trimble 1987; Carr 1994), the flat rotation curves in spiral galaxies supply the fundamental information that a large amount of dark matter is concealed in galactic halos, but unfortunately fail to provide any hint as regard to its constituents.

As first proposed by Paczyński (1986, 1996), gravitational microlensing can greatly help to clarify the nature of dark matter, and since 1993 this dream has started to become a reality with the detection of several microlensing events towards the Large Magellanic Cloud (Alcock et al. 1993; Aubourg et al. 1993).

Today, although the evidence for Massive Astrophysical Compact Halo Objects (MACHOs) – presumably located in the halo of our galaxy – is firm, the implications of such a discovery crucially depend on the assumed galactic model (Gates, Gyuk & Turner 1996)\*. What can be reliably concluded from the existing data-set is that MACHOs should lie in the mass range  $0.05 - 1.0 M_{\odot}$ , but stronger claims are unwarranted because the MACHO mass strongly depends on the uncertain properties of the considered galactic model (Evans 1996, De Paolis, Ingrosso & Jetzer 1996).

High mass values  $\gtrsim 0.5 M_{\odot}$  suggest white dwarfs as a natural candidate for MACHOs (Tamanaha et al. 1990, Adams & Laughlin 1996, Chabrier, Segretein & Méra 1996, Fields, Mathews & Schramm 1996). Besides requiring a rather *ad hoc* initial mass function sharply peaked somewhere in the range  $1 - 8 M_{\odot}$  and a halo age larger than  $\sim 16$  Gyr, this option implies an unobserved increased metallicity of the interstellar medium (Gibson & Mould 1996).

Somewhat lower mass values  $\sim 0.3 M_{\odot}$  point towards the possibility that MACHOs are hydrogen-burning main sequence M-dwarfs. However, the null results of several searches for low-mass stars in our galaxy (Hu et al. 1994) entail that the halo cannot be mostly in the form of M-dwarfs, and the same conclusion is reached by optical imaging of high-latitude fields taken with the Wide Field Camera of HST (Bahcall et al. 1994). Observe that these results are derived under the assumption of a smooth spatial distribution of M-dwarfs, and become considerably less severe in the case of a clumpy distribution (Kerins 1996, 1997).

Still lower mass values  $\lesssim 0.1 M_{\odot}$  make brown dwarfs an attractive candidate for MACHOs†. Actually, precisely the latter mass values are supported by nonstandard halo models – like the maximal disk model (van Albada & Sancisi 1986; Persic & Salucci 1990) – which tend to be favoured by recent observational data (Sackett 1996).

\* It has become customary to take the standard spherical halo model as a baseline for comparison. Unfortunately, because of the presently available limited statistics, different data-analysis procedures lead to results which are only marginally consistent. Indeed, within the standard halo model, the average MACHO mass value reported by the MACHO team is  $0.5^{+0.3}_{-0.2} M_{\odot}$  (Alcock et al. 1996), whereas the mass moment method (De Rújula, Jetzer & Massó 1991) yields  $0.27 M_{\odot}$  (Jetzer 1996).

† We stress that the limit for hydrogen burning – usually quoted as  $0.08 M_{\odot}$  – gets increased up to  $0.11 M_{\odot}$  for low-metallicity objects, such as a halo population (D’Antona 1987, Burrows, Hubbard & Lunine 1989).

In spite of the fact that present uncertainties do not permit to make any sharp statement about the nature of MACHOs, brown dwarfs nevertheless look as a viable possibility to date, and we shall stick to it throughout.

Once this idea is accepted, a few almost obvious consequences follow. Given the fact that ordinary halo stars form in (globular) clusters, it seems more likely that brown dwarfs as well form in clusters (to be referred to as *dark clusters*) rather than in isolation. Furthermore – since no known star-formation mechanism is very efficient – we expect a substantial fraction of the primordial gas to be left over. Finally – because brown dwarfs do not give rise to stellar winds – this gas should presumably remain confined within the dark clusters<sup>‡</sup>. As a matter of fact, it has been shown (De Paolis et al. 1995a, 1995b; Gerhard & Silk 1996) that these suggestions can be turned into a consistent scenario, which encompasses the Fall and Rees theory for the formation of globular clusters (Fall & Rees 1985) and indeed predicts that dark clusters of brown dwarfs and cold molecular clouds – mainly of  $H_2$  – should form in the halo at galactocentric distances larger than 10 – 20 kpc. A slightly different model based on the presence of a strong cooling-flow phase during the formation of our galaxy has been considered by Fabian and Nulsen (1994, 1997) and leads to a halo made of low-mass objects.

Before proceeding further, we would like to notice that the most promising way to test whether MACHOs are indeed clumped into dark clusters is through correlation effects in microlensing observations. For the more massive dark clusters in the range  $10^5 - 10^6 M_\odot$ , the implied degeneracy in the spatial and velocity distributions would result in a strong autocorrelation in the sky position of microlensing events on an angular scale of  $20''$ , along with a correlation in their duration. It has been pointed out that already a fairly poor statistics would be sufficient to rule out the possibility that MACHOs are clumped into such massive clusters, while to confirm it more events are needed (Maoz 1994, Metcalf & Silk 1996). On the other hand, if either the mass of the dark clusters is considerably smaller than  $10^6 M_\odot$  or the fraction of dark matter in the form of MACHOs is small, then a much larger sample of events may be required.

The aim of the present paper is to address a further issue that the above picture implies. As we shall discuss below, a large fraction of MACHOs is expected to consist of binary

<sup>‡</sup> We emphasize that a halo primarily made of *unclustered* brown dwarfs (as well as white and M-dwarfs) would contain too much diffuse hot gas (at virial temperature  $\sim 10^6$  K) which emits in the X-ray band.

brown dwarfs. However, this circumstance is an occasion for anxiety, since the overwhelming majority of MACHOs looks like unresolved objects in current microlensing experiments. We will demonstrate that such an apparent paradox does not actually exist within the model in question. For, dynamical friction on molecular clouds makes binary brown dwarfs so close that they cannot have been resolved so far – still, we argue that they can be resolved in future microlensing experiments with a more accurate photometric observation.

## 2 PRELIMINARY CONSIDERATIONS

A thorough account of the scenario under consideration has been reported elsewhere, along with an analysis of several observational tests (De Paolis et al. 1996). We should stress that the lack of observational informations about dark clusters would make any effort to understand their structure and dynamics hopeless, were it not for some remarkable insights that our unified treatment of globular and dark clusters provides us.

In the first place, it looks quite natural to assume that also dark clusters have a denser core surrounded by an extended spherical halo. Moreover, it seems reasonable to suppose (at least tentatively) that dark clusters have the same average mass density as globular clusters. Accordingly, we get  $r_{DC} \simeq 0.12 (M_{DC}/M_{\odot})^{1/3}$  pc, with  $M_{DC}$  and  $r_{DC}$  denoting the mass and median radius of a dark cluster, respectively. So, by the virial theorem the one-dimensional velocity dispersion  $\sigma$  of MACHOs and molecular clouds within a dark cluster reads

$$\sigma \simeq 6.9 \times 10^{-2} \left( \frac{M_{DC}}{M_{\odot}} \right)^{1/3} \text{ km s}^{-1} . \quad (1)$$

In addition, dark clusters – just like globular clusters – presumably stay for a long time in a quasi-stationary phase, with an average central density  $\rho(0)$  slightly lower than  $10^4 M_{\odot} \text{ pc}^{-3}$  (which is the observed average central density for globular clusters).

Below, we recall only some basic points of the considered picture, which are important for the considerations to follow. First, the amount of virialized diffuse gas inside dark clusters should be small – otherwise it should have been observed in the radio band – and so most of the gas has to be in the form of cold self-gravitating clouds clumped into dark clusters. Second – at variance with the case of globular clusters – the mass distribution of dark clusters should be smooth, with a corresponding spectrum extending from  $M_{DC} \sim 10^6 M_{\odot}$  down to much lower values. Third, the dark clusters with  $3 \times 10^2 M_{\odot} \lesssim M_{DC} \lesssim 10^6 M_{\odot}$  are expected to have survived all disruptive effects and should still populate the outer galactic halo today.

Fourth, the dark clusters with  $M_{DC} \lesssim 5 \times 10^4 M_\odot$  should have begun core collapse. Fifth – since the temperature of these clouds is close to that of the Cosmic Background Radiation (De Paolis et al. 1995c) – the virial theorem entails

$$r_m \simeq 4.8 \times 10^{-2} \left( \frac{M_m}{M_\odot} \right) \text{ pc} , \quad (2)$$

where  $r_m$  and  $M_m$  indicate the median radius and mass, respectively, of a self-gravitating cloud. Accordingly, the average number density  $n_m$  in a cloud turns out to be

$$n_m \simeq 62.2 \left( \frac{\text{pc}}{r_m} \right)^2 \text{ cm}^{-3} . \quad (3)$$

Sixth, the leftover gas is mainly  $H_2$ .

We suppose for definiteness (and with an eye to microlensing experiments) that all individual brown dwarfs have mass  $m \simeq 0.1 M_\odot$ . Furthermore, since also molecular clouds originate from the fragmentation process which produces the brown dwarfs, we suppose (for definiteness) that they lie in the mass range  $10^{-3} M_\odot \lesssim M_m \lesssim 10^{-1} M_\odot$ . Correspondingly, eqs. (2) and (3) yield  $4.8 \times 10^{-5} \text{ pc} \lesssim r_m \lesssim 4.8 \times 10^{-3} \text{ pc}$  and  $2.7 \times 10^{10} \text{ cm}^{-3} \gtrsim n_m \gtrsim 2.7 \times 10^6 \text{ cm}^{-3}$ , respectively.

### 3 MACHOS AS BINARY BROWN DWARFS

Much in the same way as it occurs for ordinary stars, also in the present case the fragmentation mechanism that gives rise to individual brown dwarfs should presumably produce a substantial fraction of binary brown dwarfs – they will be referred to as *primordial* binaries. It is important to keep in mind that their mass fraction  $f_{PB}$  can be as large as 50% and that – because of the mass stratification instability – they will concentrate inside the dark cluster cores (Spitzer 1987), which are therefore expected to be chiefly composed by binaries and molecular clouds. In addition – as far as dark clusters with  $M_{DC} \lesssim 5 \times 10^4 M_\odot$  are concerned – *tidally-captured* binary brown dwarfs ought to form in the cores because of the increased central density caused by core collapse (Fabian, Pringle & Rees 1975). Thus, we are led to the conclusion that MACHOs should consist of both *individual* and *binary* brown dwarfs within the present picture.

Our subsequent analysis will be carried out in the general case of arbitrary  $f_{PB}$ , but our interest is of course focused on the case of large  $f_{PB}$ .

We recall that a binary system is hard when its binding energy exceeds the kinetic energy of field stars (otherwise it is soft). As is well known, soft binaries always get softer whereas

hard binaries always get harder because of encounters with individual stars (Heggie 1975). In the case under consideration, binary brown dwarfs happen to be hard when their orbital radius  $a$  obeys the constraint

$$a \lesssim 1.4 \times 10^{12} \left( \frac{M_{\odot}}{M_{DC}} \right)^{2/3} \text{ km} . \quad (4)$$

Hence, only those binaries which satisfy condition (4) can survive up until the present.

Now, consistency with the results of microlensing experiments – in the sense specified above – requires that the present orbital radius of the overwhelming majority of binary brown dwarfs should be (roughly) less than one-half of the Einstein radius for microlensing towards the LMC (Gaudi & Gould 1996). This demand implies in turn that the further constraint

$$a_{\text{today}} \lesssim 3 \times 10^8 \text{ km} \quad (5)$$

has to be met today. What is crucial to realize is that the latter bound turns out to be *stronger* than the former for  $M_{DC} \lesssim 3.2 \times 10^5 M_{\odot}$ .

Let us consider first *tidally-captured* binaries, whose formation is ultimately brought about by core collapse in dark clusters and is expected to proceed as in the case of globular clusters. So, following the analysis of Press and Teukolsky (1977) and Lee and Ostriker (1986) we deduce that practically all individual brown dwarfs in a dark cluster core get captured into binaries as soon as the core density  $\S \rho(0)$  starts to satisfy the condition

$$\rho(0) \gtrsim \frac{3.2 \times 10^4}{f_{IBD}} \left( \frac{M_{DC}}{M_{\odot}} \right)^{0.4} M_{\odot} \text{ pc}^{-3} , \quad (6)$$

where  $f_{IBD}$  denotes the mass fraction of individual brown dwarfs in the core. According to the above assumptions, we expect  $\rho(0) \simeq 10^4 M_{\odot} \text{ pc}^{-3}$  *just before core collapse*, and so a moderate increase in  $\rho(0)$  may be sufficient to make tidal capture operative – hence this process would occur at the onset of core collapse. However, this conclusion depends on  $f_{PB}$ , since  $f_{IBD}$  necessarily gets small for large  $f_{PB}$ . What is the orbital radius of tidally-captured binary brown dwarfs? Following the procedure outlined by Statler, Ostriker and Cohn (1987) we easily find  $a \simeq 2.5 \times 10^5 \text{ km}$  (this value is almost independent of  $M_{DC}$ ), and so tidally-captured binaries are very hard and obviously obey condition (5). Unfortunately, tidally-captured binary brown dwarfs are irrelevant from the observational point of view, since their fractional abundance turns out to be anyway less than 1%.

$\S$  We suppose throughout that the core of dark clusters has a constant density profile.

What is the fate of *primordial* binaries? Surely, only those which are hard can survive. In fact, were individual - binary encounters the only relevant process, we would conclude that hard binary brown dwarfs should indeed survive. However, also binary - binary encounters play an important rôle in the dark cluster cores, where binaries are far more abundant than individual brown dwarfs as long as  $f_{PB}$  is not negligibly small. Now, it is well known that in the latter process one of two binaries gets often disrupted (this cannot happen for both binaries – given that they are hard – while fly bys are rather infrequent), thereby leading to the depletion of the primordial binary population. We will discuss this effect in Sect. 5, where we shall find that for realistic values of the dark cluster parameters the binary break up does not take place. Still, this is not the end of the story. For, all values for their orbital radius consistent with condition (4) are in principle allowed. Thus, it is crucial that primordial binaries manage to shrink in such a way that *also condition (5) is eventually met*.

## 4 HARDENING PROCESSES

Let us turn our attention to the specific mechanism by which condition (5) can get fulfilled. Observe that this requirement demands that a binary should give up a binding energy larger than  $4.5 \times 10^{43}$  erg.

### 4.1 Collisional hardening

Collisional hardening – namely the process whereby hard binaries get harder in encounters with individual brown dwarfs – looks as the most natural possibility. In order to see whether it works, we consider the associated average hardening rate (Spitzer & Mathieu 1980), which reads presently

$$\langle \dot{E} \rangle_{CH} \simeq - 2.8 \frac{G^2 m^3 n_{IBD}(0)}{\sigma}, \quad (7)$$

where  $n_{IBD}(0) \simeq f_{IBD} \rho(0)/m$  is the number density of individual brown dwarfs in the core. So, eq. (7) becomes

$$\begin{aligned} \langle \dot{E} \rangle_{CH} \simeq & - 1.7 \times 10^{32} f_{IBD} \left( \frac{M_{\odot}}{M_{DC}} \right)^{1/3} \times \\ & \left( \frac{\rho(0)}{M_{\odot} \text{ pc}^{-3}} \right) \text{ erg yr}^{-1}, \end{aligned} \quad (8)$$

thanks to eq. (1). From eq. (8) we can estimate the total amount of binding energy  $\Delta E_B$  released by a primordial binary via encounters with individual brown dwarfs during the

Universe lifetime ¶. Taking  $\rho(0) \sim 10^3 M_\odot \text{ pc}^{-3}$  – in agreement with our assumptions – we find  $\Delta E_B \sim 10^{43} - 10^{44} f_{IBD} \text{ erg}$  (depending on  $M_{DC}$ ). As already pointed out, a large  $f_{PB}$  would imply a small  $f_{IBD}$ , in which case collisional hardening fails to reduce the present orbital radius of primordial binary brown dwarfs down to a value consistent with condition (5). Furthermore, denoting by  $\Delta a$  the change of orbital radius corresponding to a loss of binding energy  $\Delta E_B$ , we have  $|\Delta E_B/E_B| = |\Delta a/a|$ , which shows more generally that collisional hardening turns out to be irrelevant for microlensing observations.

## 4.2 Frictional hardening

As we shall show below, the presence of molecular clouds in the dark cluster cores – which is indeed the most characteristic feature of the model in question – provides a novel mechanism by which primordial binary brown dwarfs give up a larger amount of binding energy, thereby getting so hard that condition (5) is actually obeyed. Basically, this occurs because of dynamical friction on molecular clouds, and so it will be referred to as *frictional hardening*.

It is not difficult to extend the standard treatment of dynamical friction (Binney & Tremaine 1987) to the relative motion of the brown dwarfs in a binary system which moves inside a molecular cloud. For simplicity, we assume that molecular clouds have a constant density profile  $\rho_m$ . In the case of a circular orbit ||, the equations of motion imply that the time  $t_{21}$  needed to reduce the orbital radius  $a$  from  $a_1$  down to  $a_2$  is

$$t_{21} \simeq 0.17 \left(\frac{m}{G}\right)^{1/2} \frac{1}{\rho_m \ln \Lambda} (a_2^{-3/2} - a_1^{-3/2}), \quad (9)$$

where the Coulomb logarithm reads  $\ln \Lambda \simeq \ln(r_m v_c^2 / Gm) \simeq \ln(r_m / a_1)$ , with  $v_c$  denoting the circular velocity (approximately given by Kepler’s third law). Manifestly, the diffusion approximation – upon which the present treatment is based – requires that the orbital radius of a binary should always be smaller than the median radius of a cloud. As we are concerned henceforth with hard binaries,  $a_1$  has to obey condition (4). On the other hand,  $a_1$  is the larger value for the orbital radius in eq. (9). So, we shall take for definiteness – in the Coulomb logarithm only –  $a_1 \simeq 1.4 \times 10^{12} (M_\odot / M_{DC})^{2/3} \text{ km}$ . In addition, from eq. (2) we have  $\rho_m \simeq 2.5 (\text{pc}/r_m)^2 M_\odot \text{ pc}^{-3}$ . Hence, putting everything together we obtain

$$t_{21} \simeq 5 \times 10^{25} \Xi \left(\frac{r_m}{\text{pc}}\right)^2 \left[ \left(\frac{\text{km}}{a_2}\right)^{3/2} - \left(\frac{\text{km}}{a_1}\right)^{3/2} \right] \text{ yr}, \quad (10)$$

¶ The age of the Universe is nominally taken to be  $10^{10} \text{ yr}$  throughout.

|| Indeed, the circularization of the orbit is achieved by tidal effects after a few periastron passages (Zahn 1987).

having set

$$\Xi \equiv [3 + \ln(r_m/\text{pc}) + 0.7 \ln(M_{DC}/M_\odot)]^{-1} . \quad (11)$$

Specifically, the diffusion approximation demands  $\Xi > 0$ , which yields in turn

$$r_m > 5 \times 10^{-2} \left( \frac{M_\odot}{M_{DC}} \right)^{0.7} \text{ pc} . \quad (12)$$

Observe that for  $M_{DC} \lesssim 2.1 \times 10^4 M_\odot$  this constraint restricts the range of allowed values of  $r_m$  as stated in Sect. 2.

Let us begin by estimating the time  $t_{\text{frict}}$  required by dynamical friction to reduce the orbital radius from  $a_{\text{in}} \simeq 1.4 \times 10^{12} (M_\odot/M_{DC})^{2/3}$  km down to  $a_{\text{fin}} \simeq 3 \times 10^8$  km, under the assumption that a binary moves all the time inside molecular clouds. Clearly, from eq. (10) we get

$$t_{\text{frict}} \simeq 9.5 \times 10^{12} \Xi \left( \frac{r_m}{\text{pc}} \right)^2 A \text{ yr} , \quad (13)$$

where we have set

$$A \equiv \left[ 1 - 3.1 \times 10^{-6} \left( \frac{M_{DC}}{M_\odot} \right) \right] . \quad (14)$$

As an indication, we notice that for  $r_m \simeq 10^{-3}$  pc ( $M_m \simeq 2 \times 10^{-2} M_\odot$ ) and  $M_{DC} \simeq 10^5 M_\odot$  we find  $\Xi \simeq 0.2$  and so  $t_{\text{frict}} \simeq 1.3 \times 10^6$  yr.

Were the dark clusters completely filled by clouds, eq. (13) would be the final result. However, the distribution of the clouds is lumpy. So,  $t_{\text{frict}}$  has to be understood as the total time spent by a binary inside the clouds. Thus, the total time demanded to reduce  $a_{\text{in}}$  down to  $a_{\text{fin}}$  is evidently longer than  $t_{\text{frict}}$  and can be computed by the following procedure. Keeping in mind that both the clouds and the binaries have average velocity  $v \simeq \sqrt{3}\sigma$  (for simplicity, we neglect the equipartition of kinetic energy of the binaries) it follows that the time needed by a binary to cross a single cloud is

$$t_m \simeq \frac{r_m}{\sqrt{2} v} \simeq 5.6 \times 10^6 \left( \frac{r_m}{\text{pc}} \right) \left( \frac{M_\odot}{M_{DC}} \right)^{1/3} \text{ yr} . \quad (15)$$

We see that the above values of  $r_m$  and  $M_{DC}$  entail  $t_m \simeq 1.2 \times 10^2$  yr. Therefore, frictional hardening involves many clouds and is actually accomplished only after

$$N_m \simeq \frac{t_{\text{frict}}}{t_m} \simeq 1.7 \times 10^6 \Xi \left( \frac{r_m}{\text{pc}} \right) \left( \frac{M_{DC}}{M_\odot} \right)^{1/3} A \quad (16)$$

clouds have been traversed. Taking again the above values for  $r_m$  and  $M_{DC}$ , we get  $N_m \simeq 1.1 \times 10^4$ .

It is illuminating to find how many crossings of the core are necessary for a binary to

traverse  $N_m$  clouds. To this end, we proceed to estimate the number of clouds  $N_c$  encountered during *one* crossing of the core. Describing the dark clusters by a King model, we can identify the core radius with the King radius. Evidently, the cross-section for binary - cloud encounters is  $\pi r_m^2$  and so we have

$$N_c \simeq \left( \frac{9\sigma^2}{4\pi G\rho(0)} \right)^{1/2} n_{CL}(0) \pi r_m^2, \quad (17)$$

with  $n_{CL}(0)$  denoting the cloud number density in the core. Thanks to eq. (2), we can write

$$\begin{aligned} n_{CL}(0) &\simeq f_{CL} \frac{\rho(0)}{M_m} \\ &\simeq 4.8 \times 10^{-2} f_{CL} \left( \frac{\text{pc}}{r_m} \right) \times \\ &\quad \left( \frac{\rho(0)}{M_\odot \text{ pc}^{-3}} \right) \text{ pc}^{-3}, \end{aligned} \quad (18)$$

where  $f_{CL}$  denotes the fraction of core dark matter in the form of molecular clouds. Correspondingly, eq. (17) becomes

$$\begin{aligned} N_c &\simeq 0.13 \times f_{CL} \left( \frac{\rho(0)}{M_\odot \text{ pc}^{-3}} \right)^{1/2} \times \\ &\quad \left( \frac{M_{DC}}{M_\odot} \right)^{1/3} \left( \frac{r_m}{\text{pc}} \right), \end{aligned} \quad (19)$$

on account of eq. (1). So, the total number of core crossings  $N_{cc}$  that a binary has to make in order to reduce its orbital radius down to  $a_{\text{fin}} \simeq 3 \times 10^8$  km is

$$N_{cc} \simeq \frac{N_m}{N_c} \simeq 1.3 \times 10^7 f_{CL}^{-1} \Xi \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right)^{1/2} A. \quad (20)$$

On the other hand, the core crossing time is

$$\begin{aligned} t_{cc} &\simeq \left( \frac{9\sigma^2}{4\pi G\rho(0)} \right)^{1/2} \frac{1}{v} \\ &\simeq 7 \times 10^6 \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right)^{1/2} \text{ yr}. \end{aligned} \quad (21)$$

Thus, the total time needed by frictional hardening to make primordial binary brown dwarfs unresolvable in present microlensing experiments is

$$\begin{aligned} t_{\text{tot}} &\simeq N_{cc} t_{cc} \simeq 9 \times 10^{13} f_{CL}^{-1} \times \\ &\quad \Xi \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right) A \text{ yr}. \end{aligned} \quad (22)$$

Observe that  $t_{\text{tot}}$  is insensitive to  $r_m$  and almost insensitive to  $M_{DC}$  (this dependence enters only logarithmically in  $\Xi$ ). As a matter of fact, since we have considered so far only the extreme case of largest  $a_{\text{in}}$ ,  $t_{\text{tot}}$  in eq. (22) should be understood as an *upper bound* to the total

hardening time. Accordingly – taking  $f_{CL} \simeq 0.5$  – we see that for the above illustrative values of  $r_m$  and  $M_{DC}$  we get  $t_{\text{tot}}$  shorter than the age of the Universe for  $\rho(0) \gtrsim 2.5 \times 10^3 M_\odot \text{ pc}^{-3}$ . As the latter value is consistent with our assumptions, we conclude that frictional hardening has enough time to operate.

## 5 BINARY - BINARY ENCOUNTERS

We are now in position to investigate the implications of binary - binary encounters.

As a first step, we recall that the average rate  $\Gamma$  for individual - binary (IB) and binary - binary (BB) encounters (Spitzer & Mathieu 1980) can presently be written as

$$\Gamma \simeq \alpha \frac{Gma}{\sigma}, \quad (23)$$

with  $\alpha$  being either  $\alpha_{IB} \simeq 14.3$  or  $\alpha_{BB} \simeq 13$ . Correspondingly, on account of eq. (1) the reaction time in the dark cluster cores turns out to be

$$t_{\text{react}} \simeq 10^{19} \beta \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right) \left( \frac{M_{DC}}{M_\odot} \right)^{1/3} \left( \frac{\text{km}}{a} \right) \text{ yr}, \quad (24)$$

with  $\beta$  being either  $\beta_{IB} \simeq 3.1/f_{IBD}$  or  $\beta_{BB} \simeq 6.7/f_{PB}$ . Since both processes are *a priori* of comparable strength and in the cores we expect  $f_{IBD} \ll f_{PB}$ , individual - binary encounters will be neglected.

As already stressed, binary - binary encounters can lead to their disruption. Now, if no hardening were to occur – which means that the orbital radius  $a$  would stay constant – the binary survival condition would simply follow by demanding that  $t_{\text{react}}^{BB}$  should exceed the age of the Universe. However, hardening makes  $a$  decrease, and so  $t_{\text{react}}^{BB}$  increases with time. This effect can be taken into account by considering the average value  $\langle t_{\text{react}}^{BB} \rangle$  of the reaction time over the time interval in question (to be denoted by  $T$ ), namely

$$\langle t_{\text{react}}^{BB} \rangle \equiv \frac{1}{T} \int_0^T dt t_{\text{react}}^{BB}. \quad (25)$$

In order to compute  $\langle t_{\text{react}}^{BB} \rangle$  the temporal dependence  $a(t)$  of the binary orbital radius is required. This quantity can be obtained as follows. We have seen that the total time  $t_{21}^{TOT}$  needed to reduce  $a_1$  down to  $a_2$  is actually longer than  $t_{21}$ , since the cloud distribution is lumpy. So, by going through the same steps as before, we easily find

$$t_{21}^{TOT} \simeq 4.8 \times 10^{26} \Xi f_{CL}^{-1} \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right) \times \left[ \left( \frac{\text{km}}{a_2} \right)^{3/2} - \left( \frac{\text{km}}{a_1} \right)^{3/2} \right] \text{ yr}. \quad (26)$$

Setting for notational convenience  $t \equiv t_{21}^{TOT}$ ,  $a_0 \equiv a_1$  and  $a(t) \equiv a_2$ , eq. (26) yields

$$a(t) \simeq \left[ \left( \frac{\text{km}}{a_0} \right)^{3/2} + 2.1 \times 10^{-27} \Xi^{-1} f_{CL} \times \left( \frac{\rho(0)}{M_\odot \text{pc}^{-3}} \right) \left( \frac{t}{\text{yr}} \right) \right]^{-2/3} \text{ km} . \quad (27)$$

Combining eqs. (24) and (27) together and inserting the ensuing expression into eq. (25), we get

$$\begin{aligned} < t_{react}^{BB} > \simeq 1.9 \times 10^{46} f_{PB}^{-1} f_{CL}^{-1} \Xi \left( \frac{M_{DC}}{M_\odot} \right)^{1/3} \left( \frac{\text{yr}}{T} \right) \times \\ & \left( \frac{\text{km}}{a_0} \right)^{5/2} \left( \frac{M_\odot \text{pc}^{-3}}{\rho(0)} \right)^2 \left\{ \left[ 1 + 2.1 \times 10^{-27} \Xi^{-1} f_{CL} \times \left( \frac{\rho(0)}{M_\odot \text{pc}^{-3}} \right) \left( \frac{T}{\text{yr}} \right) \left( \frac{a_0}{\text{km}} \right)^{3/2} \right]^{5/3} - 1 \right\} \text{ yr} . \end{aligned} \quad (28)$$

Let us now require  $< t_{react}^{BB} >$  to exceed the age of the Universe (taking evidently  $T \simeq 10^{10}$  yr). As it is apparent from eq. (24),  $t_{react}^{BB}$  is shorter for softer binaries. Hence, in order to contemplate *hard* binaries with an arbitrary orbital radius we have to set  $a_0 \simeq 1.4 \times 10^{12} (M_\odot / M_{DC})^{2/3}$  km in eq. (28). Correspondingly, the binary survival condition reads <sup>★★</sup>

$$\begin{aligned} f_{PB} &\lesssim 8.2 \times 10^{-5} f_{CL}^{-1} \Xi \left( \frac{M_{DC}}{M_\odot} \right)^2 \left( \frac{M_\odot \text{pc}^{-3}}{\rho(0)} \right)^2 \times \\ & \left\{ \left[ 1 + 35 f_{CL} \Xi^{-1} \left( \frac{\rho(0)}{M_\odot \text{pc}^{-3}} \right) \left( \frac{M_\odot}{M_{DC}} \right) \right]^{5/3} - 1 \right\} . \end{aligned} \quad (29)$$

Although the presence of various dark cluster parameters prevents a clear - cut conclusion to be drawn from eq. (29), in the illustrative case  $M_{DC} \simeq 10^5 M_\odot$  and  $f_{CL} \simeq 0.5$  eq. (29) entails e.g.  $f_{PB} \lesssim 0.3$  for  $\rho(0) \simeq 3 \times 10^3 M_\odot \text{pc}^{-3}$ . Thus, we infer that for realistic values of the parameters in question a sizable fraction of primordial binary brown dwarfs survives binary - binary encounters in the core <sup>††</sup>.

Binary - binary encounters also have an implication of direct relevance for frictional hardening. Indeed, before claiming that such a mechanism really accomplishes its job, we have to make sure that primordial binaries do not leave the core before their orbital radius is reduced down to acceptable values. As in the case of globular clusters, encounters between individual and binary (IB) brown dwarfs can give them enough kinetic energy to escape

<sup>★★</sup> Application of the same argument to tidally - captured binaries shows that no depletion occurs in this way.

<sup>††</sup> If the initial value of  $f_{PB}$  fails to satisfy condition (29), primordial binaries start to be destroyed in binary - binary encounters until their fractional abundance gets reduced down to a value consistent with condition (29).

from the core, and the same happens in encounters between binaries (BB) (Spitzer 1987). As explained above, we can restrict our attention to the latter process, and we consider  $\langle t_{react}^{BB} \rangle$  as averaged over the total hardening time  $t_{tot}$  (so we are going to take  $T \simeq t_{tot}$  in eq. (28)). Again, in order to contemplate *hard* binaries with an arbitrary orbital radius we set  $a_0 \simeq 1.4 \times 10^{12} (M_\odot/M_{DC})^{2/3}$  km in eq. (28). Recalling eq. (22), we find

$$\frac{\langle t_{react}^{BB} \rangle}{t_{tot}} \simeq 10^{-12} f_{CL} f_{PB}^{-1} \Xi^{-1} A^{-2} \left( \frac{M_{DC}}{M_\odot} \right) \times \left\{ \left[ 1 + 3.1 \times 10^5 A \left( \frac{M_\odot}{M_{DC}} \right) \right]^{5/3} - 1 \right\} \text{ yr} . \quad (30)$$

As in the case of eq. (29), the various parameters showing up in eq. (30) do not permit to make sharp statements, but in the illustrative case  $M_{DC} \simeq 10^5 M_\odot$ ,  $f_{CL} \simeq 0.5$  and  $f_{PB} \lesssim 0.3$  eq. (30) yields  $\langle t_{react}^{BB} \rangle \gtrsim t_{tot}$ . In conclusion, it seems fair to say that condition (5) is ultimately met by the overwhelming majority of binary brown dwarfs (this statement is especially true in view of the fact that  $t_{tot}$  – as given by eq. (22) – has to be viewed as an upper bound to the total hardening time).

## 6 ORBITAL RADIUS OF PRIMORDIAL BINARIES

Our concern was primarily addressed to show that primordial binaries are hard enough today, so as to avoid conflict with the results of microlensing experiments. Still, it would be important to know how large is their present orbital radius. Unfortunately, the lack of information about the formation mechanism prevents a clear-cut answer to be given, but it looks remarkable that a nontrivial – and in fact very interesting – lower bound can nevertheless be derived. Basically, the argument is very similar to the previous one that led us to eq. (26). However, it will now be used with a different logic, for we shall leave both  $a_1$  and  $a_2$  free while demanding that  $t_{21}^{TOT}$  should equal the age of the Universe. Correspondingly, eq. (26) implies

$$\left( \frac{\text{km}}{a_2} \right)^{3/2} \simeq \left( \frac{\text{km}}{a_1} \right)^{3/2} + 2.1 \times 10^{-17} f_{CL} \times \Xi^{-1} \left( \frac{\rho(0)}{M_\odot \text{ pc}^{-3}} \right) . \quad (31)$$

Manifestly, frictional hardening is operative to the extent that  $a_2$  becomes considerably smaller than  $a_1$ . Accordingly, from eq. (31) we see that this is indeed the case provided

$$a_1 \gtrsim 1.3 \times 10^{11} f_{CL}^{-2/3} \Xi^{2/3} \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right)^{2/3} \text{ km} . \quad (32)$$

Owing to eq. (32), eq. (31) entails

$$a_2 \simeq 1.3 \times 10^{11} f_{CL}^{-2/3} \Xi^{2/3} \left( \frac{M_\odot \text{ pc}^{-3}}{\rho(0)} \right)^{2/3} \text{ km} . \quad (33)$$

Physically, the emerging picture is as follows. Only those primordial binaries whose initial orbital radius satisfies condition (32) are affected by frictional hardening, and their present orbital radius turns out to be almost *independent* of the initial value. We can make the present discussion somewhat more specific by noticing that our assumptions strongly suggest  $\rho(0) \lesssim 10^4 M_\odot \text{ pc}^{-3}$ , in which case both eq. (32) and eq. (33) acquire the form

$$a_{1,2} \gtrsim 2.8 \times 10^8 f_{CL}^{-2/3} \Xi^{2/3} \text{ km} . \quad (34)$$

Evidently, very hard primordial binaries – which violate condition (34) – do not suffer frictional hardening, and the same is true for tidally-captured binaries. We stress that these conclusions are (practically) unaffected by collisional hardening.

## 7 ENERGY BALANCE

As the above analysis shows, dynamical friction transfers a huge amount of energy from primordial binary brown dwarfs to molecular clouds, and so it looks compelling to investigate (at least) the gross features of the energy balance.

Let us start by evaluating the energy acquired by molecular clouds in the process of frictional hardening. Recalling that the traversal time for a single cloud is given by eq. (15), eq. (10) entails that – after a binary with initial orbital radius  $a_1$  has crossed  $N$  clouds – its orbital radius gets reduced down to

$$a_{N+1} \simeq \left[ B_N \left( \frac{M_\odot}{M_{DC}} \right)^{1/3} + \left( \frac{\text{km}}{a_1} \right)^{3/2} \right]^{-2/3} \text{ km} , \quad (35)$$

with  $B_N \equiv 1.1 \times 10^{-19} N \Xi^{-1} (\text{pc}/r_m)$ . Accordingly, we see that the orbital radius remains almost constant until  $N$  reaches the critical value

$$N_* \equiv 9 \times 10^{18} \Xi \left( \frac{r_m}{\text{pc}} \right) \left( \frac{M_{DC}}{M_\odot} \right)^{1/3} \left( \frac{\text{km}}{a_1} \right)^{3/2} , \quad (36)$$

whereas it *decreases* afterwards. Because the energy acquired by the clouds is just the binding energy given up by primordial binaries, the above information can be directly used to compute the energy  $\Delta E_c(N)$  gained by the  $N$ -th cloud traversed by a binary with initial orbital radius  $a_1$ . Manifestly, we have

$$\Delta E_c(N) = \frac{1}{2} G m^2 \left( \frac{1}{a_{N+1}} - \frac{1}{a_N} \right) . \quad (37)$$

Thanks to eq. (35), a straightforward calculation shows that  $\Delta E_c(N)$  stays practically constant

$$\Delta E_c \simeq 9.8 \times 10^{32} \Xi^{-1} \left( \frac{\text{pc}}{r_m} \right) \times \left( \frac{M_\odot}{M_{DC}} \right)^{1/3} \left( \frac{a_1}{\text{km}} \right)^{1/2} \text{ erg} , \quad (38)$$

as long as  $N \lesssim N_*$ , while it subsequently *decreases*. So, the amount of energy transferred to a cloud is maximal during the early stages of hardening. Now, since the binding energy of a cloud is

$$E_c \simeq 7.7 \times 10^{42} \left( \frac{r_m}{\text{pc}} \right) \text{ erg} , \quad (39)$$

it can well happen that  $\Delta E_c > E_c$  (depending on  $r_m$ ,  $M_{DC}$  and  $a_1$ ), which means that the cloud would evaporate unless it manages to efficiently dispose of the excess energy.

A deeper insight into this issue can be gained as follows (we focus the attention on the early stages of hardening, when the effect under consideration is more dramatic). Imagine that a spherical cloud is crossed by a primordial binary which moves along a straight line, and consider the cylinder  $\Delta$  traced by the binary inside the cloud (its volume being approximately  $\pi a^2 r_m$ ). Hence, by eq. (3) the average number of molecules inside  $\Delta$  turns out to be

$$N_\Delta \simeq 5.8 \times 10^{30} \left( \frac{a_1}{\text{km}} \right)^2 \left( \frac{\text{pc}}{r_m} \right) . \quad (40)$$

Physically, the energy  $\Delta E_c$  gets first deposited within  $\Delta$  in the form of heat. Neglecting thermal conductivity (more about this, later), the temperature inside  $\Delta$  accordingly becomes

$$\begin{aligned} T_\Delta &\simeq \frac{2}{3} \frac{\Delta E_c}{N_\Delta k_B} \\ &\simeq 8.1 \times 10^{17} \Xi^{-1} \left( \frac{M_\odot}{M_{DC}} \right)^{1/3} \left( \frac{\text{km}}{a_1} \right)^{3/2} \text{ K} , \end{aligned} \quad (41)$$

$k_B$  being the Boltzmann constant. On account of eqs. (4) and (32), eq. (41) yields

$$\begin{aligned} 0.5 \Xi^{-1} \left( \frac{M_{DC}}{M_\odot} \right)^{2/3} \text{ K} &\lesssim T_\Delta \lesssim 16.2 \times \\ f_{CL} \Xi^{-2} \left( \frac{\rho(0)}{M_\odot \text{pc}^{-3}} \right) \left( \frac{M_\odot}{M_{DC}} \right)^{1/3} \text{ K} , \end{aligned} \quad (42)$$

which – in the illustrative case  $f_{CL} \simeq 0.5$ ,  $\rho(0) \simeq 3 \times 10^3 M_\odot \text{pc}^{-3}$  and  $M_{DC} \simeq 10^5 M_\odot$  – entails in turn  $5.3 \times 10^3 \text{ K} \lesssim T_\Delta \lesssim 1.3 \times 10^4 \text{ K}$ . As a consequence of the increased temperature, the molecules within  $\Delta$  will radiate, thereby reducing the excess energy in the cloud. In order to see whether this mechanism actually prevents the cloud from evaporating, we notice that the characteristic time needed to accumulate the energy  $\Delta E_c$  inside  $\Delta$  is just the traversal

time  $t_m$ . Therefore, this energy will be totally radiated away provided the cooling rate per molecule  $\Lambda$  exceeds the critical value  $\Lambda_*$  given by the equilibrium condition

$$N_\Delta \Lambda_* t_m \simeq \Delta E_c . \quad (43)$$

Specifically, eq. (43) yields

$$\Lambda_* \simeq 10^{-12} \Xi^{-1} \left( \frac{\text{pc}}{r_m} \right) \left( \frac{\text{km}}{a_1} \right)^{3/2} \text{ erg s}^{-1} \text{ mol}^{-1} , \quad (44)$$

on account of eqs. (15), (38) and (40). Moreover, in the present case in which most of the molecules are  $H_2$  the explicit form of  $\Lambda$  is (see e.g. O’Dea et al. 1994, Neufeld et al. 1995)

$$\Lambda \simeq 3.8 \times 10^{-31} \left( \frac{T_\Delta}{\text{K}} \right)^{2.9} \text{ erg s}^{-1} \text{ H}_2^{-1} , \quad (45)$$

which – thanks to eq. (41) – becomes

$$\Lambda \simeq 3.4 \times 10^{21} \Xi^{-2.9} \left( \frac{\text{km}}{a_1} \right)^{4.35} \text{ erg s}^{-1} \text{ H}_2^{-1} \quad (46)$$

(notice that  $\Lambda$  is almost independent of  $M_{DC}$ ). Now, from eqs. (44) and (46) it follows that the condition  $\Lambda \gtrsim \Lambda_*$  implies

$$a_1 \lesssim 6 \times 10^{11} \Xi^{-0.7} \left( \frac{r_m}{\text{pc}} \right)^{0.35} \text{ km} . \quad (47)$$

Again, it is difficult to figure out the relevance of condition (47) in general, but in the illustrative case of  $M_{DC} \simeq 10^5 M_\odot$  it turns out to be abundantly met for hard primordial binaries.

Thus, we conclude that the energy given up by primordial binary brown dwarfs and temporarily acquired by molecular clouds is efficiently radiated away, so that the clouds do not get dissolved by frictional hardening.

As a final comment, we stress that our estimate for  $T_\Delta$  should be understood as an upper bound, since thermal conductivity has been neglected. In addition, the above analysis implicitly relies upon the assumption  $T_\Delta < 10^4 \text{ K}$ , which ensures the survival of  $H_2$ . Actually, in spite of the fact that eq. (42) entails that this may well not be the case, our conclusion remains nevertheless true. For, higher temperatures would lead to the depletion of  $H_2$ , which correspondingly gets replaced by atomic and possibly ionized hydrogen. As is well known, in either case the resulting cooling rate would exceed the one for  $H_2$ , and so cooling would be even more efficient than estimated above (Böhringer & Hensler 1989).

## 8 CONCLUSIONS

We have shown that – within the considered model – the overwhelming majority of binary brown dwarfs is so hard today that their orbital radius is smaller than (roughly) one-half of the Einstein radius for microlensing towards LMC, thereby making them unresolvable in the microlensing experiments performed so far. Still, we have seen that not too hard primordial binaries – whose initial orbital radius obeys condition (31) – have an orbital radius today which typically turns out to be about one order of magnitude smaller than the above-mentioned Einstein radius. Therefore, we argue that not too hard primordial binary brown dwarfs can be resolved in future microlensing experiments with a more accurate photometric observation<sup>‡‡</sup>, the signature being small deviations from standard microlensing light curves (Dominik 1996). Notice that such a procedure complements the detection strategy for binaries suggested by Gaudi and Gould (1996). Finally, we point out that the mechanism considered here can naturally account for an average MACHO mass somewhat larger than  $\simeq 0.1 M_{\odot}$ .

## ACKNOWLEDGEMENTS

We would like to thank the Editor for a helpful remark. FDP has been partially supported by the Dr. Tomalla Foundation, GI by the Agenzia Spaziale Italiana and MR by the Dipartimento di Fisica Nucleare e Teorica, Università di Pavia, Pavia, Italy.

## REFERENCES

- Adams F.C., Laughlin G., 1996, *ApJ*, 468, 586
- Alcock C. et al., 1993, *Nature* 365, 621
- Alcock C. et al., 1996, *astro-ph* 9606165
- Aubourg E. et al., 1993, *Nature*, 365, 623
- Bahcall J. et al., 1994, *ApJ*, 435, L51
- Binney J., Tremaine S., 1987, *Galactic Dynamics*, Princeton University Press, Princeton
- Böhringer H., Hensler G., 1989, *A & A* 215, 147
- Burrows A., Hubbard W.B., Lunine J.I., 1989, *ApJ*, 345, 939
- Carr B., 1994, *Ann. Rev. Astron. Astrophys.*, 32, 531
- Chabrier G., Segretain L., Méra D., 1996, *ApJ*, 468, L21
- D’Antona F., 1987, *ApJ*, 320, 653
- De Paolis F., Ingrosso G., Jetzer Ph., Roncadelli M., 1995a, *Phys. Rev. Lett.*, 74, 14
- De Paolis F., Ingrosso G., Jetzer Ph., Roncadelli M., 1995b, *A & A*, 295, 567

<sup>‡‡</sup> See e.g. the ongoing experiments by the GMAN and PLANET collaborations (Proceedings of the *Second International Workshop on Gravitational Microlensing Surveys*, Orsay, 1996).

- De Paolis F., Ingrosso G., Jetzer Ph., Qadir A., Roncadelli M., 1995c, *A & A*, 299, 647
- De Paolis F., Ingrosso G., Jetzer Ph., Roncadelli M., 1996, *Int. J. Mod. Phys.*, D5, 151
- De Paolis F., Ingrosso G., Jetzer Ph., 1996, *ApJ*, 470, 493
- De Rújula A., Jetzer Ph., Massó E., 1991, *MNRAS*, 250, 348
- Dominik M., 1996, Thesis, Dortmund University
- Evans N.W., 1996, astro-ph 9611161
- Fabian A.C., Nulsen P.E.J., 1994, *MNRAS*, 269, L33
- Fabian A.C., Pringle J.E., Rees M.J., 1975, *MNRAS*, 172, 15
- Fall S.M., Rees M.J., 1985, *ApJ*, 298, 18
- Fields B.D., Mathews G., Schramm D.N., 1996, astro-ph 9603035
- Gates E.J., Gyuk G., Turner M., 1996, *Phys. Rev.*, D53, 4138
- Gaudi B.S., Gould A., 1996, astro-ph 9606104
- Gerhard O., Silk J., 1996, *ApJ*, 472, 34
- Gibson B.K., Mould J.R., 1996, astro-ph 9612133
- Heggie D.C., 1975, *MNRAS*, 173, 729
- Hu E.M. et al., 1994, *Nature*, 371, 493
- Jetzer Ph., 1996, *Helv. Phys. Acta*, 69, 179
- Kerins E.J., 1996, astro-ph 9610070
- Kerins E.J., 1997, astro-ph 9704179
- Lee H.M., Ostriker J.P., 1986, *ApJ*, 310, 176
- Maoz E., 1994, *ApJ*, 428, L5
- Metcalf R.B., Silk J., 1996, *ApJ* 464, 218
- Neufeld D.A., Lepp S., Melnick G.J., 1995, *ApJS* 100, 132
- Nulsen P.E.J., Fabian A.C., 1997, submitted
- O’Dea C.P. et al., 1994, *ApJ* 422, 467
- Paczynski B., 1986, *ApJ*, 304, 1
- Paczynski B., 1996, *Ann. Rev. Astron. Astrophys.*, 34, 419
- Persic M., Salucci P., 1990, *MNRAS*, 247, 349
- Press W.H., Teukolsky S.A., 1977, *ApJ*, 213, 183
- Sackett P.D., 1997, *ApJ*, 483, 103
- Spitzer L., 1987, *Dynamical Evolution of Globular Clusters*, Princeton University Press, Princeton
- Spitzer L., Mathieu R.D., 1980, *ApJ* 241, 618
- Statler T.S., Ostriker J.P., Cohn H.N., 1987, *ApJ*, 316, 626
- Tamanaha C.M. et al., 1990, *ApJ*, 358, 164
- Trimble V., 1987, *Ann. Rev. Astron. Astrophys.*, 25, 425
- van Albada T.S., Sancisi R. 1986, *Phil. Trans. R. Soc. London A* 320, 447
- Zahn J.P., 1987, *A & A*, 57, 383